



No. B 2435
December 2021

Self-declarations of environmental classification at Fass.se

Experiences from the reviewing process
during 2020

Authors: Lisette Graae and Kajsa Boll

Project members: Lisette Graae and Kajsa Boll



Authors: Lisette Graae and Kajsa Boll

Funded by: SIVL – the Foundation for IVL Swedish Environmental Research Institute and Lif – the Research-Based Pharmaceutical Industry

Report number B 2435

ISBN 978-91-7883-347-4

Edition Only available as PDF for individual printing

© IVL Swedish Environmental Research Institute 2021

IVL Swedish Environmental Research Institute Ltd.

P.O Box 210 60, S-100 31 Stockholm, Sweden

Phone +46-(0)10-7886500 // www.ivl.se

This report has been reviewed and approved in accordance with IVL's audited and approved management system.

Preface

IVL acts as an independent and external reviewer of environmental information that pharmaceutical companies use for environmental classification of pharmaceutical products. This report describes the experiences from the review process in the year 2020.

Table of contents

Summary	5
Sammanfattning.....	6
1 Environmental classification of pharmaceuticals at Fass.se.....	7
1.1 Background.....	7
1.2 How the classifications are made.....	8
1.3 The guideline and the reviewing process.....	8
2 Experiences from the reviewing process during 2020	9
2.1 Statistics of the review process during 2020.....	9
2.2 Development of the environmental assessment	10
3 Final results of the classification.....	11
3.1 Environmental risk assessments included in the statistics.....	11
3.2 Environmental classification of substances.....	11
3.3 Environmental risk.....	12
3.4 Potential to bioaccumulate	13
3.5 Persistence	14
4 Future outlook	15
5 Concluding remarks	15
6 References	17

Summary

Since 2005, Sweden has a unique environmental classification system for pharmaceutical substances. It is a self-declaration system where each pharmaceutical company is responsible for their own environmental information, which is published on the open web-based portal www.Fass.se. Prior to publication the environmental risk assessments are reviewed by IVL Swedish Environmental Research Institute (IVL) as an independent, external part. The present report describes the experiences from the review process during the year 2020. Data for the statistical analyses are gained both from the Fass.se database and from a spreadsheet that the audit team develops and uses to keep track of documents that have been reviewed or are under current review.

In 2020, environmental risk assessments (ERAs) were sent in for review 773 times. 68% of the reviewed assessments received the comment no remarks and were recommended to be published, whereas the other 32% were either recommended to be corrected or needed to be corrected before publication. The number of unique substances that were published at Fass.se during 2020 was 694 (which is lower than 773, because several companies may send in documents for the same substance). Of these substances 52% were exempted from classification, 27% were classified regarding environmental risk, and 21% could not gain any classification due to lack of data.

The work of improving the review system is an on-going process. As a part of this work IVL performs studies and activities to increase the knowledge of pharmaceuticals in the environment. During the last four years the focus for this type of work has been on proposing and developing a model for environmental risk assessment of pharmaceutical products. The model includes environmental risks associated with emissions of Active Pharmaceutical Ingredient (API) from production processes and carbon footprint in a life cycle perspective. It is intended to enable evaluation and comparison of environmental impacts of medically equivalent pharmaceutical products. A number of potential applications for the model were also identified, where the information may be used to reduce the environmental impact of pharmaceuticals along the value chain. In 2020 this part of the project has focused on improving the understanding of different stakeholder needs, as basis for recommendations for how the further development and implementation of the model may be performed. Three workshops and several interviews with different stakeholders were performed in order to map roles and responsibilities as well as drivers, incentives and barriers for different actors along the value chain. In September 2020 the report (B2395) "Reduce environmental impacts of pharmaceuticals along the value chain" was published. The conclusions from this study are used for recommendations regarding further development and implementation of the model.

Sammanfattning

Sedan 2005 har Sverige ett unikt system för miljöklassificering av läkemedel. Systemet bygger på självdeklaration, där varje läkemedelsföretag själva är ansvariga för miljöinformationen för sina substanser. Miljöklassificeringen publiceras på en web-baserad portal, [FASS Allmänhet - Start sida](#), som är öppen för allmänheten. Före publiceringen granskas miljödokumenten av IVL, som en oberoende extern part. Denna rapport beskriver erfarenheterna från granskningsarbetet under 2020. Data för de statistiska beräkningarna kommer dels från Fass-databasen och dels från det Excelark granskarna använder sig av för att protokollföra vilka dokument som håller på att granskas och vilka som har granskats.

Under 2020 sändes miljödokument till IVL för granskning 773 gånger. Av dessa rekommenderades 68 % direkt för publicering. De återstående 32 % blev antingen rekommenderade att genomföra en korrigering eller behövde göra en korrigering innan publicering. Antalet unika substanser som publicerades på Fass.se under 2020 var 694 (vilket är lägre än 773, eftersom flera företag kan sända in miljödokument för samma substans). 52 % av dem var undantagna klassificering, 27 % klassificerades med avseende på miljörisk och 21 % kunde inte klassificeras beroende på otillräckliga data.

Arbetet med att förbättra granskningssystemet är en pågående process. En del av detta arbete utgörs av studier och aktiviteter för att öka kunskapen om läkemedel i miljön. Under de senaste fyra åren har fokus för denna typ av arbete varit att föreslå och utveckla en modell för miljöbedömning av läkemedelsprodukter med avseende på miljörisker relaterade till utsläpp av aktiv läkemedelssubstans (API) från produktionsprocesser samt klimatavtryck i ett livscykelperspektiv. Syftet med modellen är att möjliggöra utvärdering och jämförelse av miljöpåverkan av medicinskt ekvivalenta läkemedelsprodukter. Ett antal potentiella applikationer för modellen har också identifierats, där informationen kan användas för att minska miljöpåverkan av läkemedel längs värdekedjan. Under 2020 har denna del av projektet fokuserat på att öka förståelsen för de olika intressenternas behov. Genom tre workshops och flera intervjuer har man kartlagt de olika intressenternas roller, ansvar, drivkrafter, incitament och hinder längs hela värdekedjan. Arbetet publicerades i rapporten (B2395) "Reduce environmental impacts of pharmaceuticals along the value chain" i september 2020. Slutsatserna därifrån ligger sedan till grund för rekommendationer kring hur vidare utveckling och implementering av modellen kan utföras.

1 Environmental classification of pharmaceuticals at Fass.se

1.1 Background

Pharmaceutical products are essential for health and wellbeing in our everyday life. Medicines provide enormous benefits, such as improvement in quality of life, and the demand will likely increase in the future due to a growing, ageing population, chronic/lifestyle diseases, emerging market expansion, and treatment- and technology advances. Unfortunately, benefits of the use of pharmaceuticals may come with an environmental downside. Therefore, pharmaceutical residues in the environment have become a prioritized area within environmental surveillance as well as within environmental risk assessment.

In 2005 environmental information was published at Fass.se to test a new model for classification, developed on the initiative by Lif - The Research-Based Pharmaceutical Industry in Sweden. The initiative was a response to an increasing public demand for environmental information on pharmaceuticals and an attempt to develop a model accepted both by Swedish stakeholders, but also by the global pharmaceutical industry. In 2010, environmental risk assessment had been conducted for all groups of pharmaceuticals (ATC codes) on the Swedish market.

The model was developed by a Swedish Working Group consisting of Lif, the Stockholm county council, the pharmacy chain Apoteket, the Swedish association of local authorities and regions (SKL) as well as the Swedish Medical Products Agency (MPA), in conjunction with the international pharmaceutical industry. Along with the implementation of the environmental classification system, IVL Swedish Environmental Research Institute (IVL) also runs projects with the aim to develop the system further. The Fass project is financed by Lif and the Foundation for IVL Swedish Environmental Research Institute (SIVL).

The results from the classification are being presented at Fass.se, a web based pharmaceutical portal that includes information on all approved pharmaceuticals on the Swedish market. The information is accessible not only to experts, county councils and other purchasing actors, but open to the public as well. On the Swedish market today, there are approximately 2034 active pharmaceutical ingredients (APIs) (MPA, October 2020, personal communication (S-E Hillvers and Marmar Nekoro)).

The environmental classification at Fass.se is a self-declaration system meaning that each pharmaceutical company is responsible for the environmental information published at Fass.se. Prior to publication the classifications are reviewed by IVL as an independent, external part to make sure that the classifications are based on a scientifically acceptable interpretation of the guidance for the pharmaceutical companies. The reviewing process ensures a common praxis for the implementation of the guideline among the different companies and feeds back experience from the self-declaration process to the system owners, Lif. At the same time, the review of the classifications informs the companies on the needs for the environmental risk assessments to be conducted according to the principles in the guideline (Lif 2012), in a scientifically acceptable way, thus supporting the quality and credibility of the system. The classifications are, according to the principles of the system, to be updated and reviewed every three years. In the reviewing process,

issues in need for further investigation are continuously coming up. This is due to availability of new data or knowledge in the field, as well as possibilities of comparisons across different pharmaceuticals with the same active ingredients. In order to keep its credibility, it is thus of outermost importance that the system is continuously reviewed and improved. The work on the review of the environmental risk assessments at Fass.se is conducted in close connection with related research studies, which form the bases for the development of the reviewing process.

The overall aim of the Fass-project during 2020 was to continue to develop and strengthen the Swedish environmental classification system in order to make it a powerful tool on a national level as well as to raise acceptance and interest on an international level. This included continued review of the companies' interpretation of the guideline, and in-depth discussions with Lif in cases where more guidance than the guideline contains was needed. It also included continued work with the earlier proposed model for environmental assessment of pharmaceutical products, mainly by improving the understanding of different stakeholder needs as basis for recommendations for further development and implementation of the model.

1.2 How the classifications are made

In the environmental classification of pharmaceuticals at Fass.se, the risk posed by the pharmaceuticals is differentiated in five different categories: insignificant risk, low risk, moderate risk, high risk and hazardous. In addition to the risk phrase, which concerns the risk of ecotoxicological effects, each substance is assigned hazard phrases for bioaccumulation and persistence. A substance can be exempted from classification, in accordance with the European Medicines Agency (EMA) Guideline (EMA 2006), if they are unlikely to result in significant risk to the environment, e.g., proteins, vitamins and electrolytes.

The environmental assessment at Fass.se is presented at two different levels. For the non-expert user there is a level with summary phrases describing the classifications regarding environmental risk, degradation and bioaccumulation, assigned to the substance. For the expert reader a second level includes all information that has been the basis for the self-declaration including a list of references to documents that have been used.

1.3 The guideline and the reviewing process

The guidelines to what environmental data that support and differentiate the classification steps were developed by the Lif-secretariat and the Lif Expert Group on Sustainable Development, including representatives from the industry, the Stockholm county council, the pharmacy chain Apoteket, SKL and MPA. After the deregulation of the pharmacy market in Sweden the pharmacy chain Apoteket has been replaced by the Swedish Pharmacy Association in the dialogue. The first guideline was published in 2007 (Lif 2007) and a revised document was presented in June 2012.

Before publication of environmental data at Fass.se, the risk and hazard assessments are reviewed by IVL. IVL comments on the choice of classification phrase based on the supporting data and gives recommendations to Lif whether revision is needed by the company before publication. If revision is needed, the company is encouraged to send the risk assessment for another review before publication.

The review by IVL results in comments in four categories:

- **Major deviation** – deficiencies in the submitted material leads to an inaccurate classification of risk or/and hazard and needs to be changed before publication at Fass.se
- **Minor deviation** - deficiencies in the submitted material that does not lead to an inaccurate classification of risk or/and hazard but still needs to be changed before publication at Fass.se
- **Remarks** – minor deficiencies, correction is recommended (although not mandatory) to be in full compliance with guideline
- **No remarks** – no deficiencies found in the submitted material and the document is recommended for publication.

2 Experiences from the reviewing process during 2020

2.1 Statistics of the review process during 2020

The audit team at IVL use a spreadsheet called the “progression list” as a tool to keep track of documents that have been reviewed or are under current review. The data recorded in the “progression list” is saved over time and extends back to October 2009. The statistic calculations presented here are based on data in the “progression list” for year 2020.

The total number of reviews during 2020 was 773 (taking into account that a company may send in documents for the same substance several times) and the most common assessment from IVL was to give no remarks (68%). During 2020, 694 environmental risk assessments with unique substance/pharmaceutical company-combinations were submitted for review. The highest grade of comment that each company received for their risk assessments for a specific substance is illustrated in Figure 1. The majority (92%) of the environmental risk assessments got the comments “no remarks” or “remark”. Only a minor part (8%) got a comment that needed to be corrected before publication. In total, risk assessments for 612 substances (lower than 694, since several companies may send in risk assessments for the same substance) were reviewed during 2020.




Figure 1: Distribution of the highest grade of recommendations that each company received for a specific substance that was sent in for review during 2020.

2.2 Development of the environmental assessment

Within the context as third-party reviewer, IVL also performs related studies to increase the knowledge of pharmaceuticals in the environment and to develop and improve the reviewing process.

The focus and awareness of the environmental consequences of pharmaceuticals is steadily increasing. So far much of the attention has been directed towards the environmental consequences of emissions of active pharmaceutical ingredients (API), both regarding emissions associated with residues reaching the environment subsequent to human ingestion and excrement, and emissions from production of the APIs. There is, however, a growing awareness that also other environmental aspects along the pharmaceutical life cycle should be considered, such as climate impacts, resource depletion and other local environmental impacts. A number of studies and strategies have been published highlighting the need for common models and criteria to assess, report, compare and evaluate the environmental impacts of pharmaceutical products for different applications, in order to be able to prioritize and promote actions to control and reduce the impacts.

As a response to this need of information, a work started with the objective to develop and propose a model for environmental assessment of production of pharmaceuticals, that includes two parts:

- Environmental risk assessment (ERA) of emissions of API from local production.
- Carbon footprint of pharmaceutical products in a life cycle perspective.

In the development of the proposed model, it was identified that the actual intended application of results needed to be better understood in order to prioritise and guide further development and

implementation of the model. Therefore, in mid-2019 a project started with the aim to define and evaluate needs, requirements and use of product specific environmental information by different stakeholders along the pharmaceutical value chain. This has been done through three workshops and several interviews with different stakeholders in order to map roles and responsibilities as well as drivers, incentives and barriers for different actors along the value chain. In September 2020 the report (B2395) “Reduce environmental impacts of pharmaceuticals along the value chain” was published.

3 Final results of the classification

3.1 Environmental risk assessments included in the statistics

The statistics below are based on data from a document, generated by Lif, showing a snapshot of all the risk assessments that are published at Fass.se at the time the document is generated. The data is retrieved as close to the end of every year as possible in order to ease comparisons between each year’s statistic calculations. The statistics include all the environmental risk assessments that had been published during 2020 and that could be viewed at Fass.se at 2021-01-27 (the date when the document was generated).

3.2 Environmental classification of substances

The total number of unique substances that was published at Fass.se during 2020 was 484, which corresponds to 551 environmental risk assessments. The larger number of risk assessments in comparison to the number of unique substances was due to the fact that one substance can be marketed and, thus, risk assessed by more than one company. 27% of the 484 unique substances were classified regarding environmental risk, 52% were exempted substances and another 21% were reviewed, but no classification could be made (either no data at all or not sufficient data). The distribution of the unique substances is illustrated in Figure 2, below.




Figure 2: Outcome in terms of environmental classification of substances at Fass.se (n = 484). The figure covers classification of environmental risk, i.e., not potential for degradation or bioaccumulation.

3.3 Environmental risk

The environmental risk phrase is based on the PEC/PNEC ratio (Predicted Environmental Concentration/Predicted No Effect Concentration) of the API. A classification of an insignificant risk means that the $PEC/PNEC \leq 0.1$; low risk: $0.1 < PEC/PNEC \leq 1$; moderate risk: $1 < PEC/PNEC \leq 10$ and high risk: $PEC/PNEC > 10$. When the $PEC/PNEC < 1$, but the substance is flagged as a potential PBT (Persistent, Bioaccumulative, and Toxic) or vPvB (very Persistent and very Bioaccumulative), the substance is classified as having hazardous environmental properties. Of the 129 substances (27%) classified according to environmental risk the vast majority were classified as posing an insignificant risk (80%), 10% were classified as low risk, 6% as moderate risk, 2% were classified as high risk and 2% was classified as being hazardous (Figure 3).

Two substances published during 2020 were classified as posing hazardous environmental properties: Midostaurin (medication for treatment of high-risk acute myeloid leukemia), and Regorafenib (medication for treatment of metastatic colorectal cancer, advanced gastrointestinal stromal tumours, and advanced hepatocellular carcinoma). Three substances published during 2020 were classified as posing high environmental risk: Abirateronacetate (medicine for treatment of prostate cancer), Betametason (a steroid medicine used for treatment of rheumatic disorders such as rheumatoid arthritis and systemic lupus erythematosus; skin diseases such as dermatitis and psoriasis; allergic conditions such as asthma and angioedema and more), and Levonorgestrel (oral contraceptive substance). Eight substances published during 2020 were classified as posing moderate environmental risk: Desogestrel, and Etinyloestradiol (oral contraceptive pills), Estradiol and Noretisteron (hormone replacement therapies), Etonogestrel (contraceptive implant), Ibuprofen (non-steroidal anti-inflammatory drug for treatment of pain, fever and inflammation); Paracetamol (for treatment of pain, fever and inflammation).




Figure 3: Outcome of the environmental risk assessments of pharmaceuticals at Fass.se (n = 129).

3.4 Potential to bioaccumulate

Of the 484 unique substances published at Fass.se during 2020, 198 (41%) were assessed for bioaccumulation potential. For 32 substances (7%) data to make an assessment were not available and for 254 substances (52%) a hazard phrase was not assigned. The majority of the latter were exempted substances, for which an assessment of bioaccumulation potential was not made.

As shown in Figure 4, the vast majority of the substances with a classification of the bioaccumulation potential were assigned a hazard phrase indicating a low potential to bioaccumulate (90%). For pharmaceuticals, often designed to be hydrophilic to enhance transportation in the body, this is to be expected. Many substances do also undergo metabolism to more hydrophilic forms in the human body.




Figure 4: Outcome of the classification of bioaccumulation at Fass.se (n = 198).

3.5 Persistence

Of the 484 unique substances published at Fass.se during 2020, 128 substances were classified for degradation (26%), data for classification were lacking for 102 substances (21%) and for 254 substances (52%), of which the majority were exempted substances, no hazard phrase was assigned.

In the assessment of degradability, the majority of the substances classified for degradation were assigned the phrase indicating that the substance is potentially persistent (63%) (Figure 5). Substances are classified as degradable e.g., if they have passed a ready biodegradability test (e.g., OECD 301) or sufficiently low dissipation half-lives are achieved in the OECD 308 test. Slowly degradable substances show e.g., inherent degradability (e.g., OECD 302); pass the criteria set up for the OECD 308 test or show significant biotic or abiotic degradation in other tests. However, a classification that the substance is potentially persistent does not necessarily mean that it cannot be degraded in the environment, but that lack of sufficient data result in the classification persistence or that persistence cannot be excluded. Substances within this category have failed a ready and/or inherent degradation test and/or the criteria proposed for the OECD 308 test. Substances within this category could also have been indicated to be potentially persistent, based on other standard or non-standard data.




Figure 5: Outcome of the classification of degradation at Fass.se for documents published during 2020 (n = 128).

4 Future outlook

During 2021 the Fass.se-project will continue to develop and strengthen the Swedish environmental classification system in order to make it a powerful tool on a national level and to raise acceptance and interest on an international level. The review of the companies' documents and interpretation of the guideline will continue with in depth discussions with Lif in cases where more guidance than the guideline contains is needed. During the review process the content and implementation of the guideline (Lif 2012) is continuously evaluated and discussed with in the review team at IVL and between Lif and IVL. The results of these discussions will be inputs for the updated version of the guideline that will be published during 2021. From mid-2021 the work with the development of the model for environmental assessment of pharmaceutical products will continue in a separate project.

5 Concluding remarks

- The Fass-project has now been on-going for sixteen years and has resulted in a unique collection of environmental risk assessments for pharmaceutical substances, accessible to experts, county councils and other purchasing actors, as well as the public via the web-based portal www.Fass.se.
- IVL has given feedback to Lif regarding the system as such, both from a scientific perspective as well as from a quality assurance perspective, providing possibilities to evaluate and improve the system.
- In the review of the classifications IVL has informed the companies, via Lif, on the revision needs, in order for the environmental risk assessments to be conducted according to the

principles in the guideline (Lif 2012), in a scientifically acceptable way, thus supporting the quality and credibility of the system.

- 773 risk assessments (pre-published) were checked in for review during 2020. 68% of these received no remarks and were recommended to be published; a large part of these were however substances exempted for classification. The remaining risk assessments received comments with recommendations for revisions.
- The work with improving the review process will continue with the aim to achieve a review process with no unnecessary delay in publication of the updated environmental risk assessments.
- The statistic calculations of the environmental risk assessments are based on data from a document, generated by Lif, showing a snapshot of all the risk assessments that are published at Fass.se at the time the document is generated. The statistics in this report include all the environmental risk assessments that have been published during 2020 and that could be viewed at Fass.se at 2021-01-27 (the date when the document was generated).
- Risk assessments for 612 unique substances were published at Fass.se during 2020. 27% of the unique substances (n = 129) were classified regarding environmental risk; 52% were exempted from classification and 21% were reviewed, but no classification could be made due to lack of data.
- A majority of the classified substances (80%) received the assessment insignificant risk. Two substances were classified as having hazardous environmental properties, three substances were classified with high environmental risk, and eight substances with moderate environmental risk.
- 41% of the unique substances (n = 198) were assessed for bioaccumulation potential. 90% of these were assigned a hazard phrase indicating low potential to bioaccumulate (i.e., BCF < 500 or log K_{ow} < 4, according to the Fass guideline (2012)).
- 26% of the unique substances (n = 128) were assessed for degradation. 63% of these were assigned a phrase indicating that the substance is potentially persistent.
- The work with the earlier developed and proposed model for environmental assessment of pharmaceutical products, that includes both environmental risks from emissions of API from production processes and carbon footprint in a life cycle perspective, continued. The focus was on improving the understanding of different stakeholder needs, as basis for recommendations for how the further development and implementation of the model may be performed.
- The report (B2395) “Reduce environmental impacts of pharmaceuticals along the value chain” was published in September 2020.

6 References

- European Medicines Agency (EMA) (2006). Committee for Medicinal Products for Human Use (CHMP); Guideline on the Environmental Risk Assessment of Medicinal Products for Human Use. 1 June 2006, Ref EMEA/CPMP/SWP/4447/00.
- Lif (2012). Environmental classification of pharmaceuticals at www.Fass.se. Guidance for pharmaceutical companies 2012.
- Lif (2007). Environmental classification of pharmaceuticals at www.Fass.se. Guidance for pharmaceutical companies 2007.
- Medical Products Agency (MPA), (2015). "Fördjupad analys av olika handlingsalternativ för att nå etappmålet om miljöhänsyn i läkemedelslagstiftningen inom EU och internationellt", report at <https://www.lakemedelsverket.se/sv>
- Pålsson A-C., Belleza E., Ryding S-E., Örtlund L. and Westberg E. (2019). Environmental assessment model for pharmaceutical products – Environmental risks related to Active Pharmaceutical Ingredients (API) and carbon footprint in a life cycle perspective. IVL report B2352.
- Pålsson A-C., Bjerkesjö P., and Perjo L. (2020). Reduce environmental impacts of pharmaceuticals along the value chain. Needs, requirements and use of product-specific environmental information by different actors and for different applications. IVL report B2395.



IVL Swedish Environmental Research Institute Ltd.
P.O. Box 210 60 // S-100 31 Stockholm // Sweden
Phone +46-(0)10-7886500 // www.ivl.se